



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Laboratory Calibration of Density-Dependent Lines in the EUV and Soft X-Ray Regions

J. K. Lepson, P. Beiersdorfer, M. F. Gu, P. Desai

December 9, 2010

NASA Laboratory Astrophysics Workshop
Gatlinburg, TN, United States
October 24, 2010 through October 27, 2010

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.

NASA LAW, October 25-28, 2010, Gatlinburg

Laboratory Calibration of Density-Dependent Lines in the EUV and Soft X-Ray Regions

J. K. Lepson, P. Beiersdorfer¹, M. F. Gu & P. Desai

Space Sciences Laboratory, University of California, Berkeley, CA 94720

lepson@ssl.berkeley.edu

M. Bitter & L. Roquemore

Princeton Plasma Physics Laboratory, Princeton, NJ, 08543

M. L. Reinke & J. L. Terry

Massachusetts Institute of Technology, Cambridge, MA 02139

ABSTRACT

We analyzed spectral data of Fe XXII and Ar XIV from laboratory sources in which the electron density varies by several orders of magnitude to help benchmark density-sensitive emission lines useful for astrophysics and to test the atomic models underlying the diagnostic line ratios. We found excellent agreement for Fe XXII, but poorer agreement for Ar XIV.

1. Introduction

A number of astrophysically important emission lines are sensitive to electron density in the EUV and soft X-ray regions. Lines from Fe XXII, for example, have been used in recent years as diagnostics of stellar coronae, such as the active variable AB Dor, Capella, and EX Hya (Sanz-Forcada et al. 2003, Mewe et al. 2001, Mauche et al. 2003).

Here we report spectral data of Fe XXII and Ar XIV from laboratory sources in which the electron density is known from either K-shell density diagnostics (for electron beam ion traps) or from non-spectroscopic means (tokamaks), ranging from $5 \times 10^{10} \text{ cm}^{-3}$ to $5 \times 10^{14} \text{ cm}^{-3}$. These measurements were used to test the atomic data underlying the density diagnostic line ratios, complementing earlier work (Chen et al. 2004).

¹Also Physics Division, Lawrence Livermore National Laboratory, Livermore, CA 94550

2. Methods

Spectra from the Livermore electron beam ion trap were taken with a varied-line-spacing flat-field grating spectrometer with a mean spacing of 1200 lines/mm and a CCD detector. Wavelength calibrations were periodically performed with higher orders of the well known K-Shell emission lines of N VI and N VII, commonly referred to as w and Lyman- α , respectively, and with lines of O V and O VI; see Lepson et al. (2002) for further details of spectrometer set-up and data acquisition. The electron density of the EBIT plasma was estimated via the ratio of the density-sensitive 1-2 N VI transition lines known as y and z (Chen et al. 2004).

Further measurements were taken with the same spectrometers, which were installed on the NSTX tokamak and christened “XEUS” for the lower wavelength instrument and “LoWEUS” for the higher wavelength instrument (Lepson et al. 2008, 2010). Wavelength calibrations were performed with higher orders of the C VI Lyman- α and C V w lines, as well as with O V and O VI, as on EBIT. The electron density of the plasma was measured directly with Multi-Point Thomson Scattering.

Measurements on the Alcator C-Mod tokamak at MIT were taken on a 2.2 m Rowland circle grating spectrometer with a micro-channel plate detector. Spectral resolution was lower than on EBIT and NSTX, but had time-resolution of 4–16 ms. Density was also measured with Multi-Point Thomson Scattering.

Observations of Capella (α Aurigae) were taken with the *Chandra X-Ray Observatory*’s Low Energy Transmission Grating Spectrometer. We used the minus order summed from several observations totalling ~ 500 ksec.

We used the Flexible Atomic Code (Gu 2003, 2008) to calculate emission strengths of two boron-like line pairs in iron and argon known to be sensitive to the electron density. In iron, we examined the Fe XXII line pair $(2s2p^2)_{3/2} \rightarrow (2s^22p)_{3/2}$ at 114.44 Å and $(2s2p^2)_{1/2} \rightarrow (2s^22p)_{1/2}$ at 117.17 Å, which was also used by Mewe et al. (2001) as part of an analysis of the corona of Capella. For argon we examined the Ar XIV line pair $(2s^22d)_{3/2} \rightarrow (2s^22p)_{1/2}$ at 27.47 Å and the blend $(2s^23d)_{5/2} \rightarrow (2s^22p)_{3/2}$ and $(2s^23d)_{3/2} \rightarrow (2s^22p)_{3/2}$ at 27.63 Å.

3. Results and Discussion

Spectra of Fe XXII are shown in Figures 1 (a) – 1 (d) for EBIT-II, NSTX, and Alcator C-Mod. The change in relative line strength is obvious between the low- and high-density regimes.

Figure 2 shows the line ratio curve of boron-like density-sensitive lines calculated with

the Flexible Atomic Code for Fe XXII (a) and Ar XIV (b) and the ratios measured on EBIT-II, NSTX, and Alcator C-Mod (Chen et al. 2004, Lepson et al. 2010, Reinke et al. 2010). For Fe XXII, excellent agreement is found between theory and tokamak measurements near the high density limit. Agreement at lower density is not as good, but overlaps within the error limits. We are attempting measurements around the critical density of 10^{13} cm^{-3} , where the rate of change is greatest. For Ar XIV, agreement between theory and measurements is rather poor at both low and, especially, high densities, indicating the need for improved atomic models.




Fig. 1.— Spectra of iron from plasmas of different densities: (a) XEUS spectrometer at EBIT-II, with electron density $\sim 2 \times 10^{11} \text{ cm}^{-3}$. (b) inset showing the density-sensitive lines of B-like Fe XXII. (c) LoWEUS spectrometer on NSTX tokamak, where the electron density ranged from $9 \times 10^{13} \text{ cm}^{-3}$ to $1 \times 10^{14} \text{ cm}^{-3}$. (d) Spectrum from Alcator C-Mod tokamak, where the electron density ranged from $9 \times 10^{13} \text{ cm}^{-3}$ to $5 \times 10^{14} \text{ cm}^{-3}$.

This work was supported by the NASA APRA program and the DOE General Plasma Science program. Part of this work was performed under the auspices of the U S Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344 and Princeton Plasma Physics Laboratory under Contract DE-AC02-09CH11466.

REFERENCES

- Chen, H., Beiersdorfer, P., Heeter, L. A., Liedahl, D. A., Naranjo-Rivera, K. L., Träbert, E., Gu, M. F., & Lepson, J. K., 2004, *ApJ*, 611, 598
- Gu, M. F., 2003, *ApJ*, 582, 1241




Fig. 2.— Comparison of theory with measurements: (a) Ratio curve calculated for the boron-like Fe XXII line pair 117.17 Å / 114.44 Å. (b) Ratio curve calculated for the boron-like Ar XIV line pair 27.63 Å / 27.47 Å. Solid line is calculated for a Maxwellian distribution with electron energy of 0.3 keV for tokamaks; dotted line is calculated for monoenergetic beam with electron energy of 1.0 keV for EBITs.

Gu, M. F., 2008, *Can. J. Phys.*, 86, 675

Lepson, J. K., Beiersdorfer, P., Brown, G. V., Liedahl, D. A., Utter, S. B., Brickhouse, N. S., Dupree, A. K., Kaastra, J. S., Mewe, R., & Kahn, S. M., 2002, *ApJ*, 578, 648

Lepson, J. K., Beiersdorfer, P., Bitter, M., & Kahn, S. M., 2008, *Can. J. Phys.*, 86, 175

Lepson, J. K., Beiersdorfer, P., Clementson, J., Gu, M. F., Bitter, M., Roquemore, L., Kaita, R., Cox, P. G., & Safronova, A., 2010, *J. Phys. B*, 43, 14410

Mauche, C. W., Liedahl, D. A., & Fournier, K. B., 2003, *ApJ Lett.*, 588, L101

Mewe, R., Raassen, A. J. J., Drake, J. J., Kaastra, J. S., van der Meer, R. L. J., & Porquet, D., 2001, *A&A*, 368, 888

Reinke, M. L., Beiersdorfer, P., Howard, N. T., Magee, E. W., Podpaly, Y., Rice, J. E., & Terry, J. L., 2010, *Rev. Sci. Instrum* 81, 10D736

Sanz-Forcada, J., Maggio, A., & Micela, G., 2003, *A&A*, 408, 1087